



**SK PHARMA LTD**

## **Vilofor Patient Information Leaflet**

Read this leaflet before using this product. Keep it for reference and ask your doctor or pharmacist if you have questions. Do not share a prescribed medicine with another person.

### **1 What this product contains and is used for**

Composition: Vildagliptin+Metformin 50/1000mg

Therapeutic class: DPP-4 inhibitor + biguanide. Combination glucose-lowering medicine.

Uses: Type 2 diabetes mellitus.

### **2 Before using this product**

Do not use if you are allergic to an active ingredient or any other ingredient. Get emergency help for difficulty breathing, collapse or swelling of the face, lips, tongue or throat.

Tell your doctor about kidney or liver problems, dehydration or heavy alcohol use. Ask about temporary interruption for serious illness, surgery or contrast imaging.

Other medicines: Insulin and sulfonylureas may increase low-blood-sugar risk. Other medicines affecting the kidneys need review.

Pregnancy and breastfeeding: ask a clinician before use if you are pregnant, planning pregnancy or breastfeeding. The advice depends on this product and your condition.

### **3 How to use this product**

Metformin-containing combinations are usually taken with food; confirm the schedule and release formulation on the prescription.

Missed dose: ask your pharmacist for the product-specific rule. Do not take an extra dose automatically; some medicines have special missed-dose instructions.

Too much: seek urgent medical advice or emergency care and take the container with you. Do not wait for symptoms or induce vomiting.



## **Vilofor Patient Information Leaflet**

### **4 Possible side effects**

Not everyone develops side effects. Effects listed here are a concise selection; the final product label must include the complete medicine-specific information.

Possible effects: Nausea, diarrhea, abdominal discomfort and, with prolonged metformin use, reduced vitamin B12.

When to get help: Severe weakness, vomiting, abdominal pain or unusual deep breathing may indicate lactic acidosis. Severe persistent abdominal pain may indicate pancreatitis. Seek urgent care.

Get emergency help for a severe allergic reaction. Contact your doctor or pharmacist for troublesome or persistent symptoms. Report suspected reactions to SK Pharma with the product name, batch number, dates of use and symptoms.

### **5 Storage and safe disposal**

Keep out of the sight and reach of children. Store in the original closed container and follow the temperature and light-protection conditions printed on the pack. Do not use after the expiry date or if the seal is damaged.

An expiry date on a label does not establish the period allowed after opening. Ask your pharmacist about the approved after-opening period and disposal of unused product. Do not discard medicines in household wastewater.

### **6 Presentation and ingredients**

Brand: Vilofor. Presentation: 3x10 Tablets.

Active ingredients: Vildagliptin+Metformin 50/1000mg.

Other ingredients: the full excipient list, including any sugar, alcohol, allergens, preservatives and capsule-shell components, must be completed from the approved SK Pharma formula.

### **7 Manufacturer and contact details**

SK Pharma Ltd

Salah Kakar Pharmaceutical Company

Manufacturing site: Pul-e-Charkhi Industrial Parks, Kabul, Afghanistan

Head office: Sakhi Zadah Trade Center, Jada-e-Maiwand, Kabul, Afghanistan

Telephone: +93 20 210 5610

Email: [info@skpharmas.com](mailto:info@skpharmas.com)

Webpage: [www.skpharmas.com](http://www.skpharmas.com)